

UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg

150 mg 200 mg 250 mg

For adults living with schizophrenia,

**RELAPSE PREVENTION
TREATMENT WITH UZEDY.***

INNER STRENGTH FROM YOU.

*During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY[®] once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.


IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full [Prescribing Information](#) for UZEDY, including **Boxed WARNING**.





SCHIZOPHRENIA IS COMPLEX

UZEDY MAY BE ABLE TO HELP

UZEDY® is a long-acting injectable (LAI) treatment option for schizophrenia so you don't have to take a pill every day.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.


MORE STABILITY, FEWER SYMPTOMS*



LESS RISK OF RELAPSE


UZEDY® (risperidone) extended-release injectable suspension significantly reduced the risk of relapse compared with placebo in a clinical study.†

Of the patients receiving once-monthly UZEDY, 7% relapsed vs 29% of patients receiving placebo.



FEWER SETBACKS

With once-monthly UZEDY, after 24 weeks (about 6 months) of treatment, the relapse-free rate of patients was 93% vs 72% for those receiving placebo.



GREATER STABILITY

In the clinical study, **87% of patients maintained stability** with once-monthly UZEDY; 61% of patients receiving placebo did so.

SAFETY WAS EXTENSIVELY STUDIED

The most common side effects of risperidone included slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY.


*In a clinical study of UZEDY, stability was defined based on patients who stayed outpatient, had few symptoms from a list of specific psychotic symptoms, and had symptom and clinical functioning scores below certain cutoffs for at least 4 weeks.

†During the maintenance stage of the RISE study (which varied in length of time), receiving UZEDY once monthly or once every 2 months significantly reduced the risk of relapse vs placebo.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including:

- **Uncontrolled facial or body movements (tardive dyskinesia)** that may not go away, even if you stop receiving UZEDY. Tardive dyskinesia may also start after you stop receiving UZEDY.



UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg



**FROM
THE START**

UZEDY[®] has a streamlined start—there are **no extra risperidone pills or additional starter doses** once you begin treatment.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including:

- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.
- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.

92%

of patients (58 out of 63 patients) were satisfied with UZEDY.

Patients with schizophrenia received UZEDY as part of a clinical trial and took a survey.*



SHORT NEEDLE SIZE

UZEDY uses a syringe with a short needle that injects a small amount of medicine just below the skin.

In fact, it's **one of the smallest LAI needles** used to treat schizophrenia.†



INJECTION SITE OPTIONS

UZEDY is injected just below the skin instead of deep into your muscle.

UPPER ARM

OR

ABDOMEN



FLEXIBLE DOSING

With UZEDY, you get **1 injection from your healthcare provider every 1 or 2 months.** You and your doctor will decide which option is right for you.

*Data were collected from 63 patients, 24 physicians, and 25 nurses in a prospective, cross-sectional companion survey assessing the perceptions regarding ease of use and satisfaction with UZEDY. The survey was administered after a minimum of 2 experiences prescribing, administering, or receiving UZEDY.


†UZEDY is supplied as a prefilled syringe designed to provide 0.14 to 0.7 mL of medication, depending on dose, with a 5/8-inch needle.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including:

- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.



STARTING THE CONVERSATION

Here are some questions to help you start a conversation with your healthcare provider.

- Is UZEDY® an option for me?
- How can UZEDY fit into my treatment routine?
- What are possible benefits and risks with UZEDY?
- What are the most common side effects with UZEDY?



Visit [UZEDY.com](https://www.uzedy.com) to download a full discussion guide to use at your next appointment.

IMPORTANT SAFETY INFORMATION (CONTINUED)

UZEDY may cause serious side effects, including:

- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including Boxed WARNING.

ONGOING SUPPORT

teva | Shared Solutions

Teva Shared Solutions® is a support program that can help you plan for starting and staying on UZEDY by:

- ✓ Confirming coverage and pharmacy
- ✓ Locating an injection site near you
- ✓ Giving over-the-phone nurse support

YOU COULD SAVE ON UZEDY


ELIGIBLE PATIENTS MAY PAY AS LITTLE AS \$0 PER DOSE*



To learn more about all your support options, go to UZEDY.com.

*Offer is available for patients with commercial insurance only. This offer is NOT available for patients eligible for Medicare, Medicaid, or any other form of government insurance coverage. Please see complete Terms and Conditions at www.UZEDY.com.

The Patient Savings Program for UZEDY® (risperidone) extended-release injectable suspension is available to eligible patients who have been prescribed UZEDY and have commercial prescription insurance. Eligible patients may pay as little as \$0 for once-monthly or once-every-2-month dosing options of UZEDY. Maximum program assistance per prescription and annual benefit limits per individual apply and out-of-pocket expenses may vary. Patient is responsible for costs above maximum benefit amounts. This program is restricted to residents of the United States and United States territories. Uninsured and cash-paying patients are NOT eligible for this program. Patients enrolled in any state or federally funded healthcare program, including but not limited to, Medicare, Medigap, Medicaid, VA, DOD, TRICARE, Puerto Rico Government Health Insurance Plan, Medicare-eligible patients enrolled in an employer-sponsored health plan or prescription drug benefit program for retirees, are NOT eligible for this Program. Teva Neuroscience, Inc. and its affiliates reserve the right to change, rescind, revoke, or discontinue this program at any time without notice. Please see complete Terms and Conditions at www.UZEDY.com.



APPROVED USE

UZEDY® (risperidone) extended-release injectable suspension is a prescription medicine used to treat schizophrenia in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Do not receive UZEDY if you are allergic to risperidone, paliperidone, or any of its components.

UZEDY may cause serious side effects, including:

- **Stroke in elderly people (cerebrovascular problems) that can lead to death.**
- **Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can lead to death. Seek medical attention right away** if you have any of these symptoms: high fever, severe muscle stiffness, confusion, sweating, irregular heartbeat, fast heart rate, or changes in your blood pressure.
- **Uncontrolled facial or body movements (tardive dyskinesia) that may not go away, even if you stop receiving UZEDY.** Tardive dyskinesia may also start after you stop receiving UZEDY.
- **Problems with your metabolism** that may include high blood sugar (hyperglycemia), diabetes mellitus, changes in the fat levels in your blood (dyslipidemia), and weight gain. Extremely high blood sugar can lead to coma or death. If you have diabetes or are at risk for diabetes (e.g., obesity, family history of diabetes), your healthcare provider should check your blood

sugar before you start and during treatment with UZEDY. Call your healthcare provider if you have symptoms of high blood sugar including: feeling very thirsty, hungry, sick to your stomach, weak or tired, or confused; needing to urinate more than usual; or your breath smells fruity.

- **High levels of prolactin in your blood.** UZEDY may cause a rise in the blood levels of a hormone called prolactin that may cause side effects including missed menstrual periods, decreased fertility in women, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **Decreased blood pressure (orthostatic hypotension).** You may feel lightheaded or faint when you rise too quickly from a sitting or lying position.
- **Falls.** Antipsychotic medicines like UZEDY may cause drowsiness or dizziness when you are standing, which could increase your risk for falls and related injuries.
- **Low white blood cell count.**
- **Problems thinking clearly and moving your body.** Do not drive, operate machinery, or do other dangerous activities until you know how UZEDY affects you.
- **Seizures (convulsions).**
- **Difficulty swallowing that can cause food or liquid to get into your lungs.**
- **Prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **Problems with control of your body temperature (too high or too low). Avoid getting overheated or dehydrated.**

The most common side effects of risperidone

included slow movements, stiffness, shaking, restlessness, abnormal muscle contractions or movements, drowsiness, dizziness, anxiety, blurred vision, nausea, vomiting, indigestion, diarrhea, increased saliva, constipation, dry mouth, increased appetite, weight gain, tiredness, rash, and common cold symptoms. Injection site reactions including a lump or itching were reported with UZEDY. These are not all the possible side effects of UZEDY. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

Do not drink alcohol during treatment with UZEDY.

Before receiving UZEDY, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome.
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia).
- have diabetes or have a family history of diabetes.
- have had dizziness or fainting or are being treated for high blood pressure.
- have had a low white blood cell count.
- have or have had seizures or epilepsy.
- are pregnant or plan to become pregnant during treatment with UZEDY. It is not known if UZEDY will harm your unborn baby. Use of UZEDY during the third trimester of pregnancy may cause side effects in the newborn infant, including agitation, abnormal muscle tone, tremor, drowsiness, difficulty feeding, and difficulty breathing. Seek medical attention if you notice these signs. If you become pregnant during treatment with UZEDY, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics, or call 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>.

- are breastfeeding or plan to breastfeed. If you are receiving UZEDY and are breastfeeding, monitor your infant for sleepiness, inadequate weight gain, jitteriness, tremors, and abnormal muscle movements. Seek medical care if you notice these signs.
- have or have had kidney or liver problems.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. UZEDY and other medicines may affect each other.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, **or call** 1-800-FDA-1088.

For more information about UZEDY, see the full Prescribing Information including **Boxed WARNING, or talk to your healthcare provider.**



UZEDY[®]
(risperidone) extended-release
injectable suspension

50 mg 75 mg 100 mg 125 mg
150 mg 200 mg 250 mg

ARE YOU READY FOR UZEDY?



Talk to your doctor
or visit [UZEDY.com](https://www.uzedy.com)
to learn more.

APPROVED USE

UZEDY[®] (risperidone) extended-release injectable suspension is a prescription medicine used to treat schizophrenia in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about UZEDY?

UZEDY can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). UZEDY is not approved for use in patients with dementia-related psychosis.

Please see additional Important Safety Information throughout and the full Prescribing Information for UZEDY, including **Boxed WARNING.**



© 2025 Teva Neuroscience, Inc.
RIS-41228 January 2025